


## Major Pharmaceuticals

Neomycin and Polymyxin B Sulfates and Hydrocortisone Otic Solution USP is a sterile antibacterial and anti-inflammatory solution for otic use.

Neomycin Sulfate is the sulfate salt of neomycin B and C, which are produced by the growth of *Streptomyces fradiae* Waksman (Fam. Streptomycetaceae). It has a potency equivalent of not less than 600 micrograms of neomycin standard per milligram, calculated on an anhydrous basis. The structural formulae are:


$$\begin{array}{c}
 \text{CH}_3 \\
 | \\
 \text{RCH}_2\text{CH}(\text{CH}_2)_4\text{CO}-\text{L-DAB}-\text{L-Thr}-\text{L-DAB}-\text{L-DAB} \begin{cases} \text{L-DAB-D-Phe-L-Leu} \\ \text{L-Thr-L-DAB-L-DAB} \end{cases} \cdot x\text{H}_2\text{SO}_4 \\
 | \qquad \qquad \qquad | \qquad \qquad \qquad | \qquad \qquad \qquad | \qquad \qquad \qquad | \\
 \gamma\text{-NH}_2 \qquad \qquad \gamma\text{-NH}_2 \qquad \qquad \gamma\text{-NH}_2 \qquad \qquad \gamma\text{-NH}_2 \qquad \qquad \gamma\text{-NH}_2
 \end{array}$$

Polymycin B<sub>1</sub> (R=CH<sub>3</sub>)  
 Polymycin B<sub>2</sub> (R=H)  
 DAB=α,γ - diaminobutyric acid

CC12CCC3C(C1CC2=O)C(C(C3)O)C(CO)C4=CC=CC=C4

## CLINICAL PHARMACOLOGY

Corticoids suppress the inflammatory response to a variety of agents and they may delay healing. Since corticoids may inhibit the body's defense mechanism against infection, a concomitant antimicrobial drug may be used when this inhibition is considered to be clinically significant in a particular case.

The anti-infective components in the combination are included to provide action against specific organisms susceptible to them. Neomycin sulfate and polymyxin B sulfate together are considered active against the following microorganisms: *Staphylococcus aureus*, *Escherichia coli*, *Haemophilus influenzae*, *Klebsiella-Enterobacter* species, *Neisseria* species, and *Pseudomonas aeruginosa*. This product does not provide adequate coverage against *Serratia marcescens* and streptococci, including *Streptococcus pneumoniae*. The relative potency of corticosteroids depends on the molecular structure, concentration, and release from the vehicle.

## INDICATIONS AND USAGE

For the treatment of superficial bacterial infections of the external auditory canal caused by organisms susceptible to the action of the antibiotics.

## CONTRAINDICATIONS

This product is contraindicated in those individuals who have shown hypersensitivity to any of its components. This product should not be used if the external auditory canal disorder is suspected or known to be due to cutaneous viral infection (for example, herpes simplex virus or varicella zoster virus).

## WARNINGS

Neomycin can induce permanent sensorineural hearing loss due to cochlear damage, mainly destruction of hair cells in the organ of Corti. The risk of ototoxicity is greater with prolonged use; therefore, duration of therapy should be limited to 10 consecutive days (see PRECAUTIONS: General).

Patients being treated with eardrops containing neomycin should be under close clinical observation. Due to its acidity which may cause burning and stinging, neomycin and polymyxin B sulfates and hydrocortisone otic solution should not be used in any patients with a perforated tympanic membrane.

Neomycin sulfate may cause cutaneous sensitization. A precise incidence of hypersensitivity reactions (primarily skin rash) due to topical neomycin is not known. Discontinue this product promptly if sensitization or irritation occurs.

When using neomycin-containing products to control secondary infection in the chronic dermatoses, such as chronic otitis externa or stasis dermatitis, it should be borne in mind that the skin in these conditions is more liable than is normal skin to become sensitized to many substances, including neomycin. The manifestation of sensitization to neomycin is usually a low-grade reddening with swelling, dry scaling and itching; it may be manifest simply as a failure to heal. Periodic examination for such signs is advisable, and the patient should be told to discontinue the product if they are observed. These symptoms regress quickly on withdrawing the medication.

Neomycin-containing applications should be avoided for the patient thereafter.

Contains potassium metabisulfite, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.

## PRECAUTIONS

### General

As with other antibiotic preparations, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi.

If the infection is not improved after 1 week, cultures and susceptibility tests should be repeated to verify the identity of the organism and to determine whether therapy should be changed.

Treatment should not be continued for longer than 10 days.

Allergic cross-reactions may occur which could prevent the use of any or all of the following antibiotics for the treatment of future infections: kanamycin; paromomycin; streptomycin; and possibly, gentamicin.

### Information for patients

Avoid contaminating bottle tip with material from the ear, fingers or other source. This caution is necessary if the sterility of the drops is to be preserved.

If sensitization or irritation occurs, discontinue use immediately and contact your physician.

Do not use in the eyes.

### Laboratory tests

Systemic effects of excessive levels of hydrocortisone may include a reduction in the number of circulating eosinophils and a decrease in urinary excretion of 17-hydroxycorticosteroids.

## **Carcinogenesis, mutagenesis, impairment of fertility**

Long-term studies in animals (rats, rabbits, mice) showed no evidence of carcinogenicity attributable to oral administration of corticosteroids.

## **Pregnancy**

### **Teratogenic effects**

Pregnancy Category C: Corticosteroids have been shown to be teratogenic in rabbits when applied topically at concentrations of 0.5% on days 6 to 18 of gestation and in mice when applied topically at a concentration of 15% on days 10 to 13 of gestation. There are no adequate and well-controlled studies in pregnant women. Corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

### **Nursing mothers**

Hydrocortisone appears in human milk following oral administration of the drug. Since systemic absorption of hydrocortisone may occur when applied topically, caution should be exercised when neomycin and polymyxin B sulfates and hydrocortisone otic solution is used by a nursing woman.

### **Pediatric use**

The safety and effectiveness of neomycin and polymyxin B sulfates and hydrocortisone otic solution in otitis externa have been established in the pediatric age group 2 years to 16 years of age. There is inadequate data to establish safety and effectiveness in otitis externa for pediatric patients under 2 years of age <sup>1</sup>.

---

**1**

Jones, RN, Milazzo J, Seidlin, M. Ofloxacin Otic Solution for Treatment of Otitis Externa in Children and Adults. *Arch Otolaryngol Head Neck Surg.* 1997; 123; 1193-1200.

### **Geriatric use**

Clinical studies of neomycin and polymyxin B sulfates and hydrocortisone otic solution did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

## **ADVERSE REACTIONS**

Neomycin occasionally causes skin sensitization. Ototoxicity and nephrotoxicity have also been reported (see WARNINGS). Adverse reactions have occurred with topical use of antibiotic combinations including neomycin and polymyxin B. Exact incidence figures are not available since no denominator of treated patients is available. The reaction occurring most often is allergic sensitization. In one clinical study, using a 20% neomycin patch, neomycin-induced allergic skin reactions occurred in two of 2,175 (0.09%) individuals in the general population. <sup>2</sup> In another study, the incidence was found to be approximately 1%. <sup>3</sup>

The following local adverse reactions have been reported with topical corticosteroids, especially under occlusive dressings: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, and miliaria. Stinging and burning have been reported when this product has gained access to the middle ear.

---

**2**

Leyden JJ, Kligman AM. Contact dermatitis to neomycin sulfate. *JAMA* 1979; 242: 276-1278.

**3**

Prystowsky SD, Allen AM, Smith RW, Nonomura JH, Odum RB, Akers WA. Allergic contact hypersensitivity to nickel, neomycin, ethylenediamine, and benzocaine: relationships between age, sex, history of exposure, and reactivity to standard patch tests and use tests in general population. *Arch Dermatol* 1979; 115:959-962.

## **DOSAGE AND ADMINISTRATION**

Therapy with this product should be limited to 10 consecutive days. The external auditory canal should be thoroughly cleansed and dried with a sterile cotton applicator.

For adults, four drops of the solution should be instilled into the affected ear 3 or 4 times daily. For infants and children, three drops are suggested because of the smaller capacity of the ear canal.

The patient should lie with the affected ear upward and then the drops should be instilled. This position should be maintained for 5 minutes to facilitate penetration of the drops into the ear canal. Repeat, if necessary, for the opposite ear.

If preferred, a cotton wick may be inserted into the canal and then the cotton may be saturated with the solution. This wick should be kept moist by adding further solution every 4 hours. The wick should be replaced at least once every 24 hours.

**HOW SUPPLIED**

Neomycin and Polymyxin B Sulfates and Hydrocortisone Otic Solution USP is supplied in a white plastic dropper bottle in the following size:

10 mL - Prod. No. 10009

**STERILE OTIC SOLUTION**

**Rx only**

**FOR USE IN THE EARS ONLY**

**NOT FOR USE IN THE EYES**

**STORAGE**

Store between 15°-30°C (59°- 86°F).

Securely tighten cap between uses.

**DO NOT USE IF IMPRINTED NECKBAND IS NOT INTACT**

**MANUFACTURER INFORMATION**

**Bausch & Lomb Incorporated**

Tampa, FL 33637

©Bausch & Lomb Incorporated

Revised February 2008

9072201 (Folded)

9072101 (Flat)

Prod. No. 10009

PRINCIPLE DISPLAY PANEL



**CORTOMYCIN**  
**OTIC SOLUTION**  
Neomycin and Polymyxin B  
Sulfates and Hydrocortisone  
Otic Solution USP (Sterile)

**CORTOMYCIN OTIC SOLUTION**  
Neomycin and Polymyxin B Sulfates  
and Hydrocortisone Otic Solution USP  
(Sterile)

**EACH mL CONTAINS: ACTIVES:**  
Neomycin Sulfate equivalent to 3.5 mg  
neomycin base, Polymyxin B Sulfate  
equal to 10,000 polymyxin B units,  
Hydrocortisone 10 mg (1%); **INACTIVES:**  
Propylene Glycol, Hydrochloric Acid,  
Potassium Metabisulfite, Cupric Sulfate,  
Glycerin, Purified Water. The pH range is  
2.0 to 4.5.

Manufactured by  
**Bausch & Lomb Incorporated**  
Tampa, Florida 33637



**USUAL DOSAGE:**  
See Package Insert.

**KEEP OUT OF REACH  
OF CHILDREN.**

**STORAGE:**  
Store between  
15°–30°C (59°–86°F).

NDC 0904-3141-10

**CORTOMYCIN**  
**OTIC SOLUTION**

Neomycin and Polymyxin B  
Sulfates and Hydrocortisone  
Otic Solution USP (Sterile)

**FOR USE IN THE EARS ONLY**

Rx only

**10 mL**



**DO NOT USE  
IF IMPRINTED  
NECKBAND IS  
NOT INTACT.**



0904-3141-10

Distributed by  
**Major Pharmaceuticals**  
Livonia, MI 48150 USA

**MAJOR®**  
PHARMACEUTICALS



1  
**35943**  
Vista color

XQ00071  
R.5/04-02  
CT10009



USUAL DOSAGE:  
See Package Insert.  
STORAGE: Store between  
15°-30°C (59°-86°F).  
KEEP OUT OF REACH  
OF CHILDREN.

DO NOT USE IF IMPRINTED  
NECKBOARD IS NOT INTACT.

Mfg. by Sealed Air Labs  
Tampa, Florida 33637

NDC 0904-3141-10

**CORTOMYCIN OTIC SOLUTION**  
Neomycin and Polymyxin B Sulfates  
and Hydrocortisone Otic Solution  
USP (Sterile)

Rx only

**FOR USE IN THE EAR ONLY**

10 mL

**MAJOR®**  
PHARMACEUTICALS



Distributed by Major Pharmaceuticals  
Livonia, MI 48150 USA

EACH mL CONTAINS:  
ACTIVES: Neomycin Sulfate  
equivalent to 3.5 mg  
neomycin base, Polymyxin  
B Sulfate equal to 10,000  
polymyxin B units, Hydro-  
cortisone 10 mg (1%)  
X000014 R/S/04-02 CT100000